arXiv:1401.1697v2 [math.FA] 30 Jan 2015

WEIGHTED COMPOSITION OPERATORS BETWEEN WEAK SPACES OF VECTOR-VALUED ANALYTIC FUNCTIONS

MOSTAFA HASSANLOU, JUSSI LAITILA, AND HAMID VAEZI

Abstract. We consider weighted composition operators $W_{\psi,\varphi}\colon f \mapsto \psi(f\circ\varphi)$ on spaces of analytic functions on the unit disc, which take values in some complex Banach space. We provide necessary and sufficient conditions for the boundedness and (weak) compactness of $W_{\psi,\varphi}$ on general function spaces, and in particular on weak vector-valued spaces. As an application, we characterize the weak compactness of $W_{\psi,\varphi}$ between two different vector-valued Bloch-type spaces. This result appears to be new also in the scalar-valued case.

1. Introduction

Let $\mathbb{D}$ be the open unit disc in the complex plane $\mathbb{C}$ and, for any complex Banach space X, let $H(\mathbb{D},X)$ denote the space of all analytic functions $f\colon \mathbb{D}\to X$. For two complex Banach spaces X and Y, let $L(X,Y)$ be the space of bounded linear operators from X to Y. Then a function $\psi\in H(\mathbb{D},L(X,Y))$ and an analytic self-map φ of $\mathbb{D}$ induce a linear weighted composition operator $W_{\psi,\varphi}$, which is defined from $H(\mathbb{D},X)$ to $H(\mathbb{D},Y)$, by

$$(W_{\psi,\varphi}f)(z)=\psi(z)f(\varphi(z)),\quad z\in\mathbb{D}. \tag{1.1}$$

We have $W_{\psi,\varphi}=M_\psi C_\varphi$, where M_ψ is the operator-valued multiplier $f\mapsto\psi f$ and C_φ is the composition operator $f\mapsto f\circ\varphi$. Weighted compositions appear naturally: for a large class of Banach spaces X, all linear onto isometries between X-valued H^∞ spaces are of the form (1.1) for suitable ψ and φ; see [4] and [18]. Properties of multipliers M_ψ and composition operators C_φ have been widely studied on various spaces of vector-valued analytic functions, including vector-valued Hardy, Bergman, BMOA and Bloch spaces; see, for example, [2, 12, 13, 19, 20] and the recent survey [16]. This study has further been extended to weighted composition operators; see [3, 11, 14, 21, 22, 28].

The study of composition operators on vector-valued function spaces involves some basic results which hold for large classes of function spaces. For example, if X is infinite dimensional, then composition operators are usually never compact on an X-valued analytic function space. On the other hand, for the weak compactness of C_φ it is necessary that X is reflexive and that C_φ is weakly compact on the corresponding scalar-valued function space. For the latter statement a partial converse holds for the so-called weak vector-valued function spaces: for any Banach space E of analytic functions

$f \in H(\mathbb{D}, \mathbb{C})$ such that E contains the constant functions and the closed unit ball B_E is compact in the compact open topology of $\mathbb{D}$, the weak space $wE(X)$ (modelled on E) is the space of functions $f \in H(\mathbb{D}, X)$ such that

$$\|f\|_{wE(X)} = \sup_{\|x^*\|_{X^*} \leq 1} \|x^* \circ f\|_E < \infty. \tag{1.2}$$

See [2, 16] for the above results.

The first aim of this paper is to extend such general necessary and sufficient conditions to weighted composition operators between two different function spaces, where the weight function ψ is scalar valued, i.e., $\psi \in H(\mathbb{D}, \mathbb{C})$. Consequently, we obtain characterizations of the weak compactness of weighted composition operators between many weak vector-valued spaces, such as weak Hardy and Bergman spaces.

In the second part, we apply the above results in characterizing the weak compactness of weighted composition operators between two different Bloch-type function spaces. Here the main part consists of a scalar-valued result, which also appears to be new.

Recall that an operator $T \in L(X, Y)$ is (weakly) compact if for any bounded sequence (x_n) in X there is a (weakly) convergent subsequence (Tx_{n_k}) in Y. Any compact operator is clearly weakly compact.

2. Weighted composition operators on general vector-valued function spaces

In this section we consider weighted composition operators between general spaces of vector-valued analytic functions. Most arguments are fairly straightforward extensions of known results for composition operators from earlier work (such as [19], [2] and [16]) but we present the details for completeness. Below we assume that the weight function ψ is complex valued ($\psi \in H(\mathbb{D}, \mathbb{C})$) and nonzero ($\psi(z) \neq 0$ for some $z \in \mathbb{D}$).

Suppose that A is a Banach space of analytic functions $\mathbb{D} \to \mathbb{C}$ and let $A(X)$ be an associated vector-valued Banach space of analytic functions $\mathbb{D} \to X$, where X is a complex Banach space. Assume that the following properties hold for the pair $(A, A(X))$ for all Banach spaces X (see [16, Section 2]):

- (a1) The constant maps $f(z) \equiv c$ belong to A for all $c \in \mathbb{C}$.
- (a2) $f \mapsto f \otimes x$ defines a bounded linear operator $J_x \colon A \to A(X)$ for any $x \in X$, where $(f \otimes x)(z) = f(z)x$ for $z \in \mathbb{D}$.
- (a3) $f \mapsto x^* \circ f$ defines a bounded linear operator $Q_{x^*} \colon A(X) \to A$ for any $x^* \in X^*$.
- (a4) The point evaluations δ_z, where $\delta_z(f) = f(z)$ for $f \in A(X)$, are bounded $A(X) \to X$ for all $z \in \mathbb{D}$.

It follows from these assumptions that the vector-valued constant maps $z \mapsto f_x(z) = 1 \otimes x \equiv x$ belong to $A(X)$ for all $x \in X$; see [16].

Suppose that $A(X)$ and $B(X)$ are two spaces of analytic functions $\mathbb{D} \to X$ such that the pairs $(A, A(X))$ and $(B, B(X))$ satisfy (a1) – (a4). The following result extends Corollary 2 of [16] and Proposition 1 of [2] to weighted composition operators. The result involves general operator ideals in the sense of Pietsch [24], although the remainder of the paper mainly concerns the closed ideals of compact and weakly compact operators (but see Remark 5 below). We will for clarity use the notation $\widetilde{W}_{\psi,\varphi}$ for the weighted composition operator between vector-valued spaces $A(X) \to B(X)$ and $W_{\psi,\varphi}$ for the operator between corresponding complex-valued spaces $A \to B$. We denote by I_X the identity operator of X.

Proposition 1. *Let X be a complex Banach space. Let $\psi \in H(\mathbb{D}, \mathbb{C})$ be nonzero and let φ be an analytic self-map of $\mathbb{D}$.*

1. *If $\widetilde{W}_{\psi,\varphi}$ is bounded $A(X) \to B(X)$, then $W_{\psi,\varphi}$ is bounded $A \to B$.*
2. *If $\widetilde{W}_{\psi,\varphi} \colon A(X) \to B(X)$ belongs to an operator ideal, then I_X and $W_{\psi,\varphi} \colon A \to B$ belong to the same operator ideal.*

Proof. Let $x \in X$, $x^* \in X^*$ be norm-1 vectors such that $\langle x^*, x\rangle = 1$, let $z_0 \in \mathbb{D}$ be a point such that $\psi(z_0) \neq 0$, and let j_1, j_2, j_3, j_4 be the linear operators

$$\begin{aligned} j_1 &\colon A \to A(X), && f \mapsto f \otimes x; \\ j_2 &\colon B(X) \to B, && f \mapsto x^* \circ f; \\ j_3 &\colon X \to A(X), && x \mapsto f_x; \\ j_4 &\colon B(X) \to X, && f \mapsto f(z_0)/\psi(z_0). \end{aligned}$$

By (a1) – (a4), all these operators are bounded, and

$$\begin{aligned} W_{\psi,\varphi} &= j_2 \circ \widetilde{W}_{\psi,\varphi} \circ j_1; \\ I_X &= j_4 \circ \widetilde{W}_{\psi,\varphi} \circ j_3. \end{aligned}$$

. □

The following result settles the question of compactness of $\widetilde{W}_{\psi,\varphi}$ between vector-valued spaces satisfying (a1) – (a4).

Proposition 2. *Let $\psi \in H(\mathbb{D}, \mathbb{C})$ be nonzero and let φ be an analytic self-map of $\mathbb{D}$. Then $\widetilde{W}_{\psi,\varphi}$ is compact $A(X) \to B(X)$ if and only if X is finite dimensional and $W_{\psi,\varphi}$ is compact $A \to B$.*

Proof. Proposition 1 (2) implies that if $\widetilde{W}_{\psi,\varphi}$ is compact, then X is finite dimensional and $W_{\psi,\varphi}$ is compact. The proof of the converse direction is a straightforward modification of the proof of [16, Prop. 7] and is thus omitted. □

Propositions 1 and 2 apply to a large number of function spaces, such as X-valued Hardy, Bergman, BMOA, and Bloch spaces as well as to their weak

versions (see, for example, [16] for the definitions of these function spaces). Before explicitly discussing any corollaries, let us consider the general class of weak vector-valued function spaces (1.2), introduced by Bonet, Domański and Lindström [2] in the context of composition operators. Suppose that E is a Banach space of analytic functions $f \in H(\mathbb{D}, \mathbb{C})$ satisfying the following conditions:

(b1) E contains the constant functions,
(b2) the closed unit ball B_E is compact in the compact open topology τ_{co} of $\mathbb{D}$.

Then the weak space defined in (1.2) is a Banach space; see [2]. Moreover, by [16, Lemma 13], properties (b1) and (b2) imply the assumptions (a1) – (a4), so that Propositions 1 and 2 apply to weak spaces.

The next result provides converses to Proposition 2 (1) and (2) for bounded and weakly compact weighted composition operators between two different weak spaces. It also extends [2, Proposition 11] to weighted composition operators. Recall here that the identity operator I_X of X is weakly compact if and only if X is reflexive.

Proposition 3. *Let $\psi \in H(\mathbb{D}, \mathbb{C})$ and let φ be an analytic self-map of $\mathbb{D}$. Suppose that E_1 and E_2 are spaces of analytic functions on $\mathbb{D}$ that satisfy assumptions (b1) and (b2).*

(1) *If $W_{\psi,\varphi}\colon E_1 \to E_2$ is bounded, then $\widetilde{W}_{\psi,\varphi}\colon wE_1(X) \to wE_2(X)$ is bounded.*
(2) *If X is a reflexive Banach space and $W_{\psi,\varphi}\colon E_1 \to E_2$ is compact, then $\widetilde{W}_{\psi,\varphi}$ is weakly compact $wE_1(X) \to wE_2(X)$.*

For the proof of Proposition 3 we need the following facts from [2]: The assumptions (b1) and (b2) imply that the point evaluation maps δ_z belong to E^* for each $z \in \mathbb{D}$ and that E is the dual space of V_E, where V_E is the closed linear span in E^* of $\{\delta_z \in E^*\colon z \in \mathbb{D}\}$. Moreover, by [2, Lemma 10], there is an isometric isomorphism $\chi\colon L(V_E, X) \to wE(X)$, so that

$$(\chi(T))(z) = T(\delta_z), \quad (\chi^{-1}(f))(\delta_z) = f(z), \tag{2.1}$$

hold for $T \in L(V_E, X)$, $f \in wE(X)$ and $z \in \mathbb{D}$.

Proof of Proposition 3. For (1), note that

$$\|x^* \circ (\widetilde{W}_{\psi,\varphi} f)\|_{E_2} = \|W_{\psi,\varphi}(x^* \circ f)\|_{E_2} \le \|W_{\psi,\varphi}\| \cdot \|x^* \circ f\|_{E_1},$$

so that $\|\widetilde{W}_{\psi,\varphi}\| \le \|W_{\psi,\varphi}\|$.

We next prove (2). Assume that $W_{\psi,\varphi}\colon E_1 \to E_2$ is compact. Its adjoint $(W_{\psi,\varphi})^*\colon E_2^* \to E_1^*$ satisfies

$$(W_{\psi,\varphi})^*(\delta_z) = \psi(z)\delta_{\varphi(z)}, \quad z \in \mathbb{D},$$

so that $(W_{\psi,\varphi})^*(V_{E_2}) \subset V_{E_1}$. We obtain the factorization $\widetilde{W}_{\psi,\varphi} = \chi_2 \circ S_{\psi,\varphi} \circ \chi_1^{-1}$, where $S_{\psi,\varphi}$ is the operator composition map

$$T \mapsto I_{L(V_{E_1},X)} \circ T \circ (W_{\psi,\varphi})^*|_{V_{E_2}}; \quad L(V_{E_2},X) \to L(V_{E_1},X),$$

and $\chi_1 \colon L(V_{E_1},X) \to wE_1(X)$ and $\chi_2 \colon L(V_{E_2},X) \to wE_2(X)$ are isometric isomorphisms from (2.1). Indeed,

$$\begin{aligned}((\chi_2 \circ S_{\psi,\varphi} \circ \chi_1^{-1})(f))(z) &= (\chi_2(\chi_1^{-1}(f) \circ (W_{\psi,\varphi})^*|_{V_{E_2}})(z) \\ &= \chi_1^{-1}(f)((W_{\psi,\varphi})^*(\delta_z)) \\ &= \chi_1^{-1}(f)(\psi(z)\delta_{\varphi(z)}) \\ &= \psi(z)\chi_1^{-1}(f)(\delta_{\varphi(z)}) \\ &= \psi(z)(f(\varphi(z)),\end{aligned}$$

for $f \in E_1$ and $z \in \mathbb{D}$.

Since $(W_{\psi,\varphi})^*|_{V_{E_2}}$ is a compact operator $V_{E_2} \to V_{E_1}$ by duality, and I_X is weakly compact by reflexivity of X, it follows from [25, Theorem 2.9] (and [26, Remark 2.4]), that the operator composition $S_{\psi,\varphi}$ is weakly compact $L(V_{E_2},X) \to L(V_{E_1},X)$. Consequently $\widetilde{W}_{\psi,\varphi}$ is weakly compact $wE_1(X) \to wE_2(X)$. □

As concrete examples, let us next consider weak versions of two classical function spaces: the Hardy space H^1 and the Bergman space A^1 (see, e.g., [29] for the definitions of H^1 and A^1). Both of these spaces satisfy (a2) and (b2), so the associated weak spaces $wH^1(X)$ and $wA^1(X)$ are Banach spaces for all X. However, they are quite different from the usual "strong" spaces $H^1(X)$ and $A^1(X)$; see [12, 13, 15].

Compactness and weak compactness of $W_{\psi,\varphi}$ are well understood on H^1 and A^1: Function-theoretic characterizations exist for the compactness of $W_{\psi,\varphi}$ on both H^1 and A^1 [7, 9]. In [7] it was further shown that every weakly compact $W_{\psi,\varphi}$ on H^1 is compact. Moreover, because A^1 is isomorphic to ℓ^1, it has the Schur property, and so every weakly compact linear operator on A^1 is compact; see [19, p. 302]. Using Proposition 3, we get the following extension of these scalar results.

Corollary 4. *Let $\psi \in H(\mathbb{D},\mathbb{C})$ be nonzero and let φ be an analytic self-map of $\mathbb{D}$. Then*

1. *$\widetilde{W}_{\psi,\varphi}$ is weakly compact on $wH^1(X)$ if and only if X is reflexive and $W_{\psi,\varphi}$ is compact on H^1.*
2. *$\widetilde{W}_{\psi,\varphi}$ is weakly compact on $wA^1(X)$ if and only if X is reflexive and $W_{\psi,\varphi}$ is compact on A^1.*

Corollary 4 could include more examples. For example, (weak) compactness of $W_{\psi,\varphi}$ between weighted spaces H^∞_v is well known [5, 6]. Also the weak space $wH^\infty_v(X)$ is well defined (and coincides with the usual "strong"

space $H_v^\infty(X)$). However, weak compactness of the more general "operator-weighted" compositions $\widetilde{W}_{\psi,\varphi}$ (where $\psi \in H(\mathbb{D}, L(X,Y))$) between $H_v^\infty(X)$ spaces has been characterized in [14].

It is worthwhile to note that the general framework (Propositions 1, 2 and 3) does not in general hold for operator-weighted composition operators. For example, an operator-weighted composition between weighted $H^\infty(X)$ spaces can be compact even if X is infinite dimensional [14]. Indeed, the present knowledge of operator-valued composition opeators is quite rudimentary and, for example, their boundedness and compactness on Hardy spaces $H^p(X)$ is still an open question; see the discussion in [16, Section 7].

Remark 5. An analogue of Proposition 3 (2) holds for the ideal of weakly conditionally compact operators. Recall that an operator $T \in L(X,Y)$ is weakly conditionally compact if for any bounded sequence (x_n) in X there is a weakly Cauchy subsequence (Tx_{n_k}) in Y. Then, if X does not contain an isomorphic copy of ℓ^1 and $W_{\psi,\varphi}\colon E_1 \to E_2$ is compact, it follows that $\widetilde{W}_{\psi,\varphi}$ is weakly conditionally compact $wE_1(X) \to wE_2(X)$. The proof is analogous to Proposition 3 (2), but uses [17] instead to deduce that the operator composition $S_{\psi,\varphi}$ is weakly conditionally compact.

3. Weighted composition operators between Bloch-type spaces

In this section we apply the results from the previous section to weighted composition operators between Bloch-type spaces of analytic functions. For $\alpha \in (0,\infty)$ and a complex Banach space X, the vector-valued Bloch-type space $\mathcal{B}_\alpha(X)$ is the Banach space of functions $f \in H(\mathbb{D}, X)$ such that

$$\|f\|_{\mathcal{B}_\alpha(X)} = \|f(0)\|_X + \sup_{z\in\mathbb{D}} \|f'(z)\|_X (1-|z|^2)^\alpha < \infty. \tag{3.1}$$

In the special case $X = \mathbb{C}$, we get the usual scalar-valued Bloch-type spaces $\mathcal{B}_\alpha = \mathcal{B}_\alpha(X)$. In the case $\alpha = 1$, we get the Bloch space $\mathcal{B}(X) = \mathcal{B}_1(X)$ (see, for example, [1]).

Boundedness and compactness of weighted composition operators between scalar-valued Bloch-type spaces $\mathcal{B}_\alpha$ is well known. Ohno, Stroethoff and Zhao [23] provided the following function-theoretic characterizations, which involve the quantities

$$q_1(\alpha,\beta,z) = \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha}|\psi(z)||\varphi'(z)|;$$

$$q_2(\beta,z) = (1-|z|^2)^\beta \log\frac{1}{1-|\varphi(z)|^2}|\psi'(z)|;$$

$$q_3(\alpha,\beta,z) = \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}|\psi'(z)|,$$

for $\alpha,\beta \in (0,\infty)$:

(1) If $0 < \alpha < 1$, then $W_{\psi,\varphi}\colon \mathcal{B}_\alpha \to \mathcal{B}_\beta$ is bounded (respectively compact) if and only if

$$\sup_{z\in\mathbb{D}} q_1(\alpha,\beta,z) < \infty$$

(respectively

$$\lim_{|\varphi(z)|\to 1} q_1(\alpha,\beta,z) = 0).$$

(2) If $\alpha = 1$, then $W_{\psi,\varphi}\colon \mathcal{B} \to \mathcal{B}_\beta$ is bounded (respectively compact) if and only if

$$\sup_{z\in\mathbb{D}} q_1(1,\beta,z) < \infty \quad \text{and} \quad \sup_{z\in\mathbb{D}} q_2(\beta,z) < \infty$$

(respectively

$$\lim_{|\varphi(z)|\to 1} q_1(1,\beta,z) = 0 \quad \text{and} \quad \lim_{|\varphi(z)|\to 1} q_2(\beta,z) = 0).$$

(3) If $1 < \alpha < \infty$, then $W_{\psi,\varphi}\colon \mathcal{B}_\alpha \to \mathcal{B}_\beta$ is bounded (respectively compact) if and only if

$$\sup_{z\in\mathbb{D}} q_1(\alpha,\beta,z) < \infty \quad \text{and} \quad \sup_{z\in\mathbb{D}} q_3(\alpha,\beta,z) < \infty$$

(respectively

$$\lim_{|\varphi(z)|\to 1} q_1(\alpha,\beta,z) = 0 \quad \text{and} \quad \lim_{|\varphi(z)|\to 1} q_3(\alpha,\beta,z) = 0).$$

The following theorem extends the scalar results to vector-valued Bloch-type spaces. For composition operators, the case $\alpha = \beta = 1$ is from [19, Thm. 4] and the case $\alpha = \beta \in (0,\infty)$ follows from [2, Prop. 11]. Our proof applies a similar approach to that of [19].

Theorem 6. *Let $\psi \in H(\mathbb{D},\mathbb{C})$ be nonzero and let φ be an analytic self-map of $\mathbb{D}$. Let $\alpha,\beta \in (0,\infty)$. Let X be a complex Banach space.*

(1) *$W_{\psi,\varphi}$ is weakly compact $\mathcal{B}_\alpha \to \mathcal{B}_\beta$ if and only if it is compact.*
(2) *$\widetilde{W}_{\psi,\varphi}$ is bounded $\mathcal{B}_\alpha(X) \to \mathcal{B}_\beta(X)$ if and only if $W_{\psi,\varphi}$ is bounded $\mathcal{B}_\alpha \to \mathcal{B}_\beta$.*
(3) *$\widetilde{W}_{\psi,\varphi}$ is weakly compact $\mathcal{B}_\alpha(X) \to \mathcal{B}_\beta(X)$ if and only if $W_{\psi,\varphi}$ is (weakly) compact $\mathcal{B}_\alpha \to \mathcal{B}_\beta$ and X is reflexive.*

For the proof of Theorem 6, we will need the following facts. For $\alpha \in (0,\infty)$, let $\mathcal{B}_{\alpha,0}$ denote the little Bloch-type space, which is the closed subspace of $\mathcal{B}_\alpha$ for which

$$\lim_{|z|\to 1} |f'(z)|(1-|z|^2)^\alpha = 0.$$

First, by [30, Thm. 15], the Bergman space A^1 is the dual space of $\mathcal{B}_{\alpha,0}$ for all α under the pairing

$$\langle f,g\rangle_\alpha = \lim_{r\to 1}\int_{\mathbb{D}} f(rz)\overline{g(rz)}(1-|z|^2)^{\alpha-1}\,\mathrm{d}A(z), \quad f \in \mathcal{B}_{\alpha,0}, g \in A^1, \quad (3.2)$$

where $\mathrm{d}A(z)$ is the normalized area measure on $\mathbb{D}$. Second, each $\mathcal{B}_{\alpha,0}$ has the Dunford-Pettis property (DPP). Recall that a Banach space E has the DPP whenever every weakly compact operator from E to some Banach space is completely continuous, i.e., maps weakly null sequences to norm null sequences. The fact that $\mathcal{B}_{\alpha,0}$ has the DPP follows from the above duality and the fact that $A^1 \sim \ell^1$ has the Schur property; see, for example, [10, 19].

Proof of Theorem 6. It is easy to see that each $\mathcal{B}_\alpha(X)$ is a weak space of the form (1.2). In fact, $\mathcal{B}_\alpha(X) = w\mathcal{B}_\alpha(X)$, with equivalent norms, where $w\mathcal{B}_\alpha(X)$ is modelled on $\mathcal{B}_\alpha$. Therefore parts (2) and (3) of Theorem 6 follow from part (1) and Propositions 1 and 3. We thus only need to prove part (1).

Let first $\alpha, \beta \in (0,\infty)$ and suppose that $W_{\psi,\varphi}$ is weakly compact, but not compact, $\mathcal{B}_\alpha \to \mathcal{B}_\beta$. Then there are $\varepsilon > 0$ and a sequence $(z_n) \subset \mathbb{D}$ so that $|\varphi(z_n)| \to 1$ and $q_1(\alpha,\beta,z_n) \geq \varepsilon$ for all n. For $n \geq 0$, define the functions $f_n \in H(\mathbb{D},\mathbb{C})$ by

$$f_n(z) = \frac{(1-|\varphi(z_n)|^2)^2}{(1-\overline{\varphi(z_n)}z)^{\alpha+1}} - \frac{1-|\varphi(z_n)|^2}{(1-\overline{\varphi(z_n)}z)^{\alpha}},$$

for $z \in \mathbb{D}$. In [23, p. 202] it has been shown that the sequence (f_n) is uniformly bounded in $\mathcal{B}_\alpha$, $f_n(z) \to 0$ uniformly on compact subsets of $\mathbb{D}$, $f_n(\varphi(z_n)) = 0$, and $f_n'(\varphi(z_n)) = \overline{\varphi(z_n)}/(1-|\varphi(z_n)|^2)^\alpha$. Thus

$$\begin{aligned} \|W_{\psi,\varphi}f_n\|_{\mathcal{B}_\beta} &\geq |(W_{\psi,\varphi}f_n)'(z_n)|(1-|z_n|)^\beta \\ &= |\varphi(z_n)|q_1(\alpha,\beta,z_n) \\ &\geq \varepsilon/2, \end{aligned} \tag{3.3}$$

for all n large enough. We further have $f_n \in \mathcal{B}_{\alpha,0}$ for all n. Let $\widehat{f_n}(j)$ denote the jth Taylor coefficient of f_n and let $p_N(z) = \sum_{j=0}^N a_j z^j$ be a polynomial. Since $f_n(z) \to 0$ uniformly on compact subsets of $\mathbb{D}$, we have $\widehat{f_n}(j) = f_n^{(j)}(0)/j! \to 0$ as $n \to \infty$. By (3.2),

$$\begin{aligned} \langle f_n, p_N\rangle_\alpha &= \lim_{r\to 1}\int_0^{2\pi}\int_0^1 \sum_{k=0}^\infty\sum_{j=0}^N \widehat{f_n}(k)\overline{a_j}r^{k+j}s^{k+j+1}\mathrm{e}^{i(k-j)\theta}(1-s^2)^{\alpha-1}\,\mathrm{d}s\frac{\mathrm{d}\theta}{2\pi} \\ &= \sum_{j=0}^N \widehat{f_n}(j)\overline{a_j}\int_0^1 s^{2j+1}(1-s^2)^{\alpha-1}\,\mathrm{d}s \to 0, \end{aligned}$$

as $n \to \infty$. Since the polynomials are dense in A^1, this means that $f_n \to 0$ weakly in $\mathcal{B}_{\alpha,0}$. Weak compactness of $W_{\psi,\varphi}$ and the DPP of $\mathcal{B}_{\alpha,0}$ now imply that $W_{\psi,\varphi}f_n \to 0$ in $\mathcal{B}_{\beta,0}$ and thus in $\mathcal{B}_\beta$. This contradicts (3.3).

We next divide the proof into 3 cases: (i) $\alpha \in (0,1)$, (ii) $\alpha = 1$, and (iii) $\alpha \in (1,\infty)$. The above argument proves the case (i). Cases (ii) and (iii) are

completed similarly, by using functions

$$g_n(z) = \frac{-1}{\log(1-|\varphi(z_n)|^2)}\left(3\left(\log\frac{1}{1-\overline{\varphi(z_n)}z}\right)^2 - 2\left(\log\frac{1}{1-\overline{\varphi(z_n)}z}\right)^3\right),$$

respectively,

$$h_n(z) = (\alpha+1)\frac{1-|\varphi(z_n)|^2}{(1-\overline{\varphi(z_n)}z)^\alpha} - \alpha\frac{(1-|\varphi(z_n)|^2)^2}{(1-\overline{\varphi(z_n)}z)^{\alpha+1}},$$

instead of f_n. □

Acknowledgement

We thank Hans-Olav Tylli for his comments on a preliminary version of the manuscript.

References

[1] J. Arregui and O. Blasco, *Bergman and Bloch spaces of vector-valued functions*, Math. Nachr. **261/262** (2003), 3–22.

[2] J. Bonet, P. Domański and M. Lindström, *Weakly compact composition operators on analytic vector-valued function spaces*, Ann. Acad. Sci. Fenn. Math. **26** (2001), 233–248.

[3] J. Bonet, M.C. Gómez-Collado, D. Jornet and E. Wolf, *Operator-weighted composition operators between weighted spaces of vector-valued analytic functions*, Ann. Acad. Sci. Fenn. Math. **37** (2012), 319–338.

[4] M. Cambern and K. Jarosz, *Multipliers and isometries in H^∞_E*, Bull. London Math. Soc. **22** (1990), 463–466.

[5] M.D. Contreras and S. Díaz-Madrigal, *Compact-type operators defined on H^∞*, Contemp. Math. **232** (1999), 111–118.

[6] M.D. Contreras and A.G. Hernández-Díaz, *Weighted composition operators in weighted Banach spaces of analytic functions*, J. Austral. Math. Soc. **69** (2000), 41–60.

[7] M.D. Contreras and A.G. Hernández-Díaz, *Weighted composition operators on Hardy spaces*, J. Math. Anal. Appl. **263** (2001), 224–233.

[8] C.C. Cowen and B.D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, CRC Press, Boca Raton 1995.

[9] Ž. Čučković and R. Zhao, *Weighted composition operators on the Bergman space*, J. London Math. Soc. **70** (2004), 499–511.

[10] J. Diestel, *A survey of results related to the Dunford-Pettis property*, Contemp. Math. **2** (1980), 15–60.

[11] M. Hassanlou, H. Vaezi and M. Wang, *Weighted composition operators on weak vector-valued weighted Bergman spaces and Hardy spaces*, Banach J. Math. Anal. **9** (2015), 35–43.

[12] J. Laitila, *Weakly compact composition operators on vector-valued BMOA*, J. Math. Anal. Appl. **308** (2005), 730–745.

[13] J. Laitila and H.-O. Tylli, *Composition operators on vector-valued harmonic functions and Cauchy transforms*, Indiana Univ. Math. J. **55** (2006), 719–746.

[14] J. Laitila and H.-O. Tylli, *Operator-weighted composition operators on vector-valued analytic function spaces*, Illinois J. Math. **53** (2009), 1019–1032.

[15] J. Laitila, H.-O. Tylli and M. Wang, *Composition operators from weak to strong spaces of vector-valued analytic functions*, J. Operator Theory **62** (2009), 281–295.
[16] J. Laitila and H.-O. Tylli, *Composition operators on vector-valued analytic function spaces: a survey*, Acta et Comm. Univ. Tartu. Math. **18** (2014), 135–157.
[17] M. Lindström and G. Schlüchtermann, *Composition of operator ideals*, Math. Scand. **84** (1999), 284–296.
[18] P.-K. Lin, *The isometries of $H^\infty(E)$*, Pacific J. Math. **143** (1990), 69–77.
[19] P. Liu, E. Saksman and H.-O. Tylli, *Small composition operators on analytic vector-valued function spaces*, Pacific J. Math. **184** (1998), 295–309.
[20] J.S. Manhas, *Multiplication operators on weighted locally convex spaces of vector-valued analytic functions*, Southeast Asian Bull. Math. **27** (2003), 649–660.
[21] J.S. Manhas, *Weighted composition operators on weighted spaces of vector-valued analytic functions*, J. Korean Math. Soc. **45** (2008), 1203–1220.
[22] J.S. Manhas, *Weighted composition operators between weighted spaces of vector-valued holomorphic functions on Banach spaces*, Appl. Math. Comput. **218** (2011), 929–934.
[23] S. Ohno, K. Stroethoff, and R. Zhao, *Weighted composition operators between Bloch-type spaces*, Rocky Mountain J. Math. **33** (2003),191–215.
[24] A. Pietsch, *Operator Ideals*, North-Holland, 1980.
[25] E. Saksman and H.-O. Tylli, *Weak compactness of multiplication operators on spaces of bounded linear operators*, Math. Scand. **70** (1992), 91–111.
[26] E. Saksman and H.-O. Tylli, *Multiplications and elementary operators in the Banach space setting*, in: Methods in Banach space theory, London Math. Soc. Lecture Note Ser. vol. 337 (2006), Cambridge University Press, Cambridge, pp. 253-292.
[27] J.H. Shapiro, *Composition Operators and Classical Function Theory*, Springer-Verlag, New York, 1993.
[28] M. Wang, *Weighted composition operators between Dirichlet spaces*, Acta Math. Sci. Ser. B Engl. Ed. **31** (2011), 641–651.
[29] K. Zhu, *Operator Theory in Function Spaces*, Marcel Dekker, New York, 1990.
[30] K. Zhu, *Bloch type spaces of analytic functions*, Rocky Mountain J. Math. **23** (1993), 1143–1177.

M. Hassanlou, Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran
E-mail address: m_hasanloo@tabrizu.ac.ir

J. Laitila, Department of Biosciences, P.O. Box 65, FI-00014 University of Helsinki, Helsinki, Finland
E-mail address: jussi.laitila@helsinki.fi

H. Vaezi, Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran
E-mail address: hvaezi@tabrizu.ac.ir